

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
10 June 2004 (10.06.2004)

PCT

(10) International Publication Number  
**WO 2004/048378 A1**

(51) International Patent Classification<sup>7</sup>: **C07D 471/06**,  
A61K 31/437, A61P 37/02

(21) International Application Number:  
PCT/SE2003/001805

(22) International Filing Date:  
21 November 2003 (21.11.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
0203471-8 22 November 2002 (22.11.2002) SE  
60/428240 22 November 2002 (22.11.2002) US  
0301299-4 6 May 2003 (06.05.2003) SE  
0301851-2 25 June 2003 (25.06.2003) SE  
60/482122 25 June 2003 (25.06.2003) US

(71) Applicant (for all designated States except US): **ACTIVE BIOTECH AB** [SE/SE]; Scheelevägen 22, Box 724, S-220 07 Lund (SE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **MATTHEWS, Ian, Richard** [GB/GB]; c/o Avidex, 57C Milton Park, GB-Abingdon, Oxfordshire OX14 4RX (GB). **COULTER, Thomas, Stephen** [GB/GB]; Evotec OAI, 111 Milton Park, Abingdon, Oxon OX14 4SD (GB). **GHIRON, Chiara** [GB/GB]; Evotec OAI, 151, Milton Park, Abingdon, Oxfordshire OX14 4SD (GB). **BRENNAN, Chris, James** [GB/GB]; Evotec OAI, 151, Milton Park, Abingdon, Oxfordshire OX14 4SD (GB). **UDDIN, Muhammed**,

**Kamal** [GB/GB]; Evotec OAI, 151, Milton Park, Abingdon, Oxfordshire OX14 4SD (GB). **PETTERSSON, Lars, Olof, Göran** [SE/SE]; c/o Active Biotech AB, Scheelevägen 22, S-220 07 Lund (SE). **THRIGE, Dortha da Graça** [DK/SE]; c/o Active Biotech AB, Scheelevägen 22, S-220 07 Lund (SE). **HUXLEY, Philip** [GB/GB]; 57c Milton Park, Abingdon, Oxfordshire OX14 4RX (GB).

(74) Agent: **AWAPATENT AB**; Box 5117, S-200 71 Malmö (SE).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ (utility model), DE (utility model), DK (utility model), DK, DM, DZ, EC, EE (utility model), EE, EG, ES, FI (utility model), FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PYRAZOLOQUINOLINES WITH IMMUNOMODULATING ACTIVITY

(57) Abstract: The present invention relates to novel heterocyclic compounds, to methods for their preparation, to compositions containing them, and to methods and use for clinical treatment of medical conditions which may benefit from immunomodulation, including rheumatoid arthritis, multiple sclerosis, diabetes, asthma, transplantation, systemic lupus erythematosus and psoriasis. More particularly the present invention relates to novel heterocyclic compounds, which are CD80 antagonists capable of inhibiting the interactions between CD80 and CD28.



WO 2004/048378 A1

### Pyrazoloquinolines with immunomodulating activity


The present invention relates to novel heterocyclic compounds, to methods for their preparation, to compositions containing them, and to methods and use for clinical treatment of medical conditions which may benefit  
5 from immunomodulation, including rheumatoid arthritis, multiple sclerosis, diabetes, asthma, transplantation, systemic lupus erythematosus and psoriasis. More particularly the present invention relates to novel heterocyclic compounds, which are CD80 antagonists capable of  
10 inhibiting the interactions between CD80 and CD28.

#### Background of the invention

The immune system possesses the ability to control the homeostasis between the activation and inactivation of lymphocytes through various regulatory mechanisms during and after an immune response. Among these are mechanisms that specifically inhibit and/or turn off an immune  
15 response. Thus, when an antigen is presented by MHC molecules to the T-cell receptor, the T-cells become properly activated only in the presence of additional co-stimulatory signals. In the absence of accessory signals there is no lymphocyte activation and either a state of functional inactivation termed anergy or tolerance is induced, or the T-cell is specifically deleted by apoptosis. One such co-stimulatory signal involves interaction of  
20 CD80 on specialised antigen-presenting cells with CD28 on T-cells, which has been demonstrated to be essential for full T-cell activation. (Lenschow et al. (1996) *Annu. Rev. Immunol.*, 14, 233-258)


A paper by Erbe et al, in *J. Biol. Chem.* Vol. 277, No. 9, pp 7363-7368 (2002), describes three small  
30 molecule ligands which bind to CD80, and inhibit binding of CD80 to CD28 and CTLA4. Two of the disclosed ligands are fused pyrazolones of structures A and B:

2



# DESCRIPTION OF THE INVENTION

According to the present invention there is provided a compound of formula (I) or a pharmaceutically or veterinarily acceptable salt thereof:



wherein

25  $R_1$  and  $R_3$  independently represent H; F; Cl; Br;  $-\text{NO}_2$ ;  $-\text{CN}$ ;  $\text{C}_1\text{-C}_6$  alkyl optionally substituted by F or Cl; or  $\text{C}_1\text{-C}_6$  alkoxy optionally substituted by F;

$R_2$  represents H, or optionally substituted  $\text{C}_1\text{-C}_6$  alkyl,  $\text{C}_3\text{-C}_7$  cycloalkyl or optionally substituted phenyl;

30 Y represents  $-\text{O}-$ ,  $-\text{S}-$ , N-oxide, or  $-\text{N}(\text{R}_5)-$  wherein  $\text{R}_5$  represents H or  $\text{C}_1\text{-C}_6$  alkyl;

X represents a bond or a divalent  $\text{C}_1\text{-C}_6$  alkylene radical;

$R_4$  represents  $-\text{C}(=\text{O})\text{NR}_6\text{R}_7$ ,  $-\text{NR}_7\text{C}(=\text{O})\text{R}_6$ ,  $-\text{NR}_7\text{C}(=\text{O})\text{OR}_6$ ,  $-\text{NHC}(=\text{O})\text{NHR}_6$ , or  $-\text{NHC}(=\text{S})\text{NHR}_6$  wherein

35  $R_6$  represents H, or a radical of formula  $-(\text{Alk})_b\text{-Q}$  wherein b is 0 or 1, and

Alk is an optionally substituted divalent straight chain or branched C<sub>1</sub>-C<sub>12</sub> alkylene, C<sub>2</sub>-C<sub>12</sub> alkenylene or C<sub>2</sub>-C<sub>12</sub> alkynylene radical which may be interrupted by one or more non-adjacent -O-, -S- or -N(R<sub>8</sub>)- radicals wherein R<sub>8</sub> represents H or C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>3</sub>-C<sub>4</sub> alkenyl, C<sub>3</sub>-C<sub>4</sub> alkynyl, or C<sub>3</sub>-C<sub>6</sub> cycloalkyl, and

Q represents H; -CF<sub>3</sub>; -OH; -SH; -NR<sub>8</sub>R<sub>8</sub> wherein each R<sub>8</sub> may be the same or different; an ester group; or an optionally substituted phenyl, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, C<sub>5</sub>-C<sub>7</sub> cycloalkenyl or heterocyclic ring having from 5 to 8 ring atoms; and

R<sub>7</sub> represents H or C<sub>1</sub>-C<sub>6</sub> alkyl; or when taken together with the atom or atoms to which they are attached R<sub>6</sub> and R<sub>7</sub> form an optionally substituted heterocyclic ring having from 5 to 8 ring atoms.

Compounds of general formula (I) are CD80 antagonists. They inhibit the interaction between CD80 and CD28 and thus the activation of T cells, thereby modulating the immune response.

Accordingly the invention also includes:

(i) a compound of formula (I) or a pharmaceutically or veterinarily acceptable salt thereof for use in the treatment of conditions which benefit from immunomodulation.

(ii) the use of a compound of formula (I) or a pharmaceutically or veterinarily acceptable salt thereof in the manufacture of a medicament for the treatment of conditions which benefit from immunomodulation,.

(iii) a method of immunomodulation in humans and non-human primates, comprising administration to a subject in need of such treatment an immunomodulatory effective dose of a compound of formula (I) or a pharmaceutically or veterinarily acceptable salt thereof.

(iv) a pharmaceutical or veterinary composition comprising a compound of formula (I) or a pharmaceutically or veterinarily acceptable salt thereof together with a

pharmaceutically or veterinarily acceptable excipient or carrier.

Conditions which benefit from immunomodulation include:

- 5 Adrenal insufficiency
- Allergic angiitis and granulomatosis
- Amyloidosis
- Ankylosing spondylitis
- Asthma
- 10 Autoimmune Addison's disease
- Autoimmune alopecia
- Autoimmune chronic active hepatitis
- Autoimmune hemolytic anemia
- Autoimmune neutropenia
- 15 Autoimmune thrombocytopenic purpura
- Autoimmune vasculitides
- Behçet's disease
- Cerebellar degeneration
- Chronic active hepatitis
- 20 Chronic inflammatory demyelinating polyradiculoneuropathy
- Dermatitis herpetiformis
- Diabetes
- Eaton-Lambert myasthenic syndrome
- Encephalomyelitis
- 25 Epidermolysis bullosa
- Erythema nodosa
- Gluten-sensitive enteropathy
- Goodpasture's syndrome
- Graft versus host disease
- 30 Guillain-Barre syndrome
- Hashimoto's thyroiditis
- Hyperthyroidism
- Idiopathic hemochromatosis
- Idiopathic membranous glomerulonephritis
- 35 Minimal change renal disease
- Mixed connective tissue disease
- Multifocal motor neuropathy

- Multiple sclerosis
- Myasthenia gravis
- Opsoclonus-myoclonus syndrome
- Pemphigoid
- 5 Pemphigus
- Pernicious anemia
- Polyarteritis nodosa
- Polymyositis/dermatomyositis
- Post-infective arthritides
- 10 Primary biliary sclerosis
- Psoriasis
- Reactive arthritides
- Reiter's disease
- Retinopathy
- 15 Rheumatoid arthritis
- Sclerosing cholangitis
- Sjögren's syndrome
- Stiff-man syndrome
- Subacute thyroiditis
- 20 Systemic lupus erythematosus
- Systemic sclerosis (scleroderma)
- Temporal arteritis
- Thromboangiitis obliterans
- Transplantation rejection
- 25 Type I and type II autoimmune polyglandular syndrome
- Ulcerative colitis
- Uveitis
- Wegener's granulomatosis

As used herein the term "alkylene" refers to a  
 30 straight or branched alkyl chain having two unsatisfied  
 valencies, for example  $-\text{CH}_2-$ ,  $-\text{CH}_2\text{CH}_2-$ ,  $-\text{CH}_2\text{CH}_2\text{CH}_2-$ ,  
 $-\text{CH}(\text{CH}_3)\text{CH}_2-$ ,  $-\text{CH}(\text{CH}_2\text{CH}_3)\text{CH}_2\text{CH}_2\text{CH}_3$ , and  $-\text{C}(\text{CH}_3)_3$ .

As used herein the term "heteroaryl" refers to a 5-  
 or 6- membered aromatic ring containing one or more he-  
 35 teroatoms. Illustrative of such groups are thienyl, fu-  
 ryl, pyrrolyl, imidazolyl, benzimidazolyl, thiazolyl,  
 pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, thia-

diazolyl, oxadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl.

As used herein the unqualified term "heterocyclyl" or "heterocyclic" includes "heteroaryl" as defined above, and in particular means a 5-8 membered aromatic or non-aromatic heterocyclic ring containing one or more heteroatoms selected from S, N and O, including for example, pyrrolyl, furanyl, thienyl, piperidinyl, imidazolyl, oxazolyl, isoxazolyl, thiazolyl, thiadiazolyl, pyrazolyl, pyridinyl, pyrrolidinyl, pyrimidinyl, morpholinyl, piperazinyl, indolyl, morpholinyl, benzofuranyl, pyranyl, isoxazolyl, quinuclidinyl, aza-bicyclo[3.2.1]octanyl, benzimidazolyl, methylenedioxyphenyl, maleimido and succinimido groups.

Unless otherwise specified in the context in which it occurs, the term "substituted" as applied to any moiety herein means substituted with one or more of the following substituents, namely (C<sub>1</sub>-C<sub>6</sub>)alkyl, trifluoromethyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy (including the special case where a ring is substituted on adjacent ring C atoms by methylenedioxy or ethylenedioxy), trifluoromethoxy, (C<sub>1</sub>-C<sub>6</sub>)alkylthio, phenyl, benzyl, phenoxy, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, hydroxy, mercapto, amino, fluoro, chloro, bromo, cyano, nitro, oxo, -COOH, -SO<sub>2</sub>OH, -CONH<sub>2</sub>, -SO<sub>2</sub>NH<sub>2</sub>, -COR<sup>A</sup>, -COOR<sup>A</sup>, -SO<sub>2</sub>OR<sup>A</sup>, -NHCOR<sup>A</sup>, -NHSO<sub>2</sub>R<sup>A</sup>, -CONHR<sup>A</sup>, -SO<sub>2</sub>NHR<sup>A</sup>, -NHR<sup>A</sup>, -NR<sup>A</sup>R<sup>B</sup>, -CONR<sup>A</sup>R<sup>B</sup> or -SO<sub>2</sub>NR<sup>A</sup>R<sup>B</sup> wherein R<sup>A</sup> and R<sup>B</sup> are independently a (C<sub>1</sub>-C<sub>6</sub>)alkyl group. In the case where "substituted" means substituted by (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, phenyl, benzyl or phenoxy, the ring thereof may itself be substituted with any of the foregoing, except (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl phenyl, benzyl or phenoxy.

As used herein the unqualified term "carbocyclyl" or "carbocyclic" refers to a 5-8 membered ring whose ring atoms are all carbon.

Some compounds of the invention contain one or more chiral centres because of the presence of asymmetric carbon atoms. The presence of asymmetric carbon atoms gives

rise to stereoisomers or diastereoisomers with R or S stereochemistry at each chiral centre. The invention includes all such stereoisomers and diastereoisomers and mixtures thereof.

5           Salts of salt forming compounds of the invention include physiologically acceptable acid addition salts for example hydrochlorides, hydrobromides, sulphates, methane sulphonates, p-toluenesulphonates, phosphates, acetates, citrates, succinates, lactates, tartrates, fumarates and maleates; and base addition salts, for  
10           example sodium, potassium, magnesium, and calcium salts. Where the compound contains an amino group, quaternary amino salts are also feasible, and are included in the invention.

15           In the compounds of the invention the following are examples of the several structural variables:

$R_1$  may be, for example, H, F, Cl, methyl, methoxy, or methylenedioxy. Currently it is preferred that  $R_1$  is H, Cl or especially F;

20            $R_2$  may be, for example H, methyl, methoxy, cyclopropyl, phenyl, or fluoro-, chloro-, methyl, or methoxy-substituted phenyl. H or cyclopropyl is presently preferred;

$R_3$  may be, for example, H, F, Cl, methyl, methoxy,  
25           or methylenedioxy. Currently it is preferred that  $R_3$  is F or Cl, and it is most preferred that  $R_3$  be H;

          Y may be, for example, -O-, -S-, or -N( $R_5$ )- wherein  $R_5$  represents H or methyl. -NH- or -S- is presently preferred.

30           X may be, for example a bond, or a -CH<sub>2</sub>- or -CH<sub>2</sub>CH<sub>2</sub>- radical. A bond is presently preferred.

$R_4$  represents -C(=O)NR<sub>6</sub>R<sub>7</sub>, -NR<sub>7</sub>C(=O)R<sub>6</sub>, -NR<sub>7</sub>C(=O)OR<sub>6</sub>, -NHC(=O)NHR<sub>6</sub>, or -NHC(=S)NHR<sub>6</sub>. Of these -NR<sub>7</sub>C(=O)R<sub>6</sub>, and especially -C(=O)NR<sub>6</sub>R<sub>7</sub> and -NHC(=O)NHR<sub>6</sub> are currently  
35           preferred.  $R_7$  is preferably H, but a wide range of  $R_6$  substituents have given rise to highly active compounds



of the invention. Many exemplary  $R_6$  substituents appear in the compounds of the Examples below.

$R_6$  may be, for example, H or a radical of formula  $-\text{Alk}_b\text{-Q}$  wherein b is 0 or 1 and

5        Alk may be, for example a  $-(\text{CH}_2)_n-$ ,  
 $-\text{CH}((\text{CH}_2)_m\text{CH}_3)(\text{CH}_2)_n-$ ,  $-\text{C}((\text{CH}_2)_m\text{CH}_3)((\text{CH}_2)_p\text{CH}_3)(\text{CH}_2)_n-$ ,  
 $-(\text{CH}_2)_n\text{-O-}(\text{CH}_2)_m-$ ,  $-(\text{CH}_2)_n\text{-NH-}(\text{CH}_2)_m-$ , or  
 $-(\text{CH}_2)_n\text{-NH-}(\text{CH}_2)_m\text{-NH-}(\text{CH}_2)_p-$  radical where n is 1, 2, 3 or  
 4 and m and p are independently 0, 1, 2, 3 or 4, and

10        Q may represent H,  $-\text{OH}$ ,  $-\text{COOCH}_3$ , phenyl, cyclopropyl, cyclopentyl, cyclohexyl, pyridyl, furyl, thienyl, or oxazolyl; and



$R_7$  may be, for example, H, or when taken together with the atom or atoms to which they are attached  $R_6$  and

15         $R_7$  may form a heterocyclic ring of 5, 6 or 7 members.

Specific examples of  $R_4$  groups include those present in the compounds of the Examples herein.


Compounds of the invention may be prepared by synthetic methods known in the literature, from compounds  
 20        which are commercially available or are accessible from commercially available compounds. For example, compounds of formula (I) wherein  $R_4$  is a group  $-\text{NR}_7\text{C(=O)R}_6$  may be prepared by acylation of an amine of formula (II) with an acid chloride of formula (III):

25




9

Compounds of the invention wherein  $R_4$  is a group  $-NHC(=O)NHR_6$  may be prepared by reaction of an amine of formula (IIA) with an isocyanate of formula (IIIA)



5

Compounds of the invention wherein  $R_4$  is a group  $-C(=O)NHR_6$  may be prepared by reaction of an acid chloride of formula (IIB) with an amine  $NHR_6R_7$ :



10

Compounds of the invention wherein  $R_4$  is a group  $-NR_7C(=O)OR_6$  may be prepared by reaction of an amine of formula (II) with a chloroformate  $ClC(=O)OR_6$ .


10

The following Examples illustrate the preparation of compounds of the invention:

Preparation of Intermediate 1

2-(4-Nitrophenyl)-6-fluoro-2,5-dihydropyrazolo[4,3-c]-  
5 quinolin-3-one

10




4-Nitrophenylhydrazine (2.28 g, 0.014 mol) was added in one portion to a stirred solution of 4-chloro-8-fluoro-quinoline-3-carboxylic acid ethyl ester (3.58 g, 0.014 mol) in anhydrous n-butyl alcohol (50 ml) at room temperature. The mixture was refluxed for 16 h under nitrogen, cooled to room temperature and then filtered to leave an orange solid. The solid was purified by washing sequentially with ethyl acetate (20 ml) and heptane (20 ml) and then finally dried under suction to give the pyrazolone (3.93 g, 87 %) as a dark orange solid, LCMS m/z 325.24 [M+H]<sup>+</sup> @ R<sub>T</sub> 1.47 min.

Preparation of Intermediate 2

25 2-(4-Aminophenyl)-6-fluoro-2,5-dihydropyrazolo[4,3-c]-  
quinolin-3-one

30




Tin (II) chloride dihydrate (12.5 g, 0.055 mol) was added in one portion to a stirred solution of 2-(4-nitrophenyl)-6-fluoro-2,5-dihydro-pyrazolo[4,3-c]quinolin-3-one (intermediate 1) (3.59 g, 0.011 mol) in ethyl alcohol (110 ml) at room temperature. The mixture was then heated

to 80 °C for 8 h, cooled to room temperature and filtered to leave a yellow solid. The solid was suspended in a bi-phasic solution of ethyl acetate (1L), a saturated solution of Rochelles salt (500 ml) and a saturated solution of sodium bicarbonate (500 ml) and stirred at room temperature for 2h. The mixture was filtered and the remaining solid was washed with water and dried under vacuum to afford the title compound (3.39 g, 99 %) as a bright yellow solid, LCMS m/z 295.30 [M+H]<sup>+</sup> @ R<sub>T</sub> 0.84 min.

10 Example 1

*N*-[4-(6-Fluoro-3-oxo-3,5-dihydropyrazolo[4,3-*c*]quinolin-2-yl)-phenyl]-2-methyl-butyramide



20 (±)-2-Methylbutyryl chloride (13.6 µl, 0.11 mmol) was added dropwise over 30 sec to a stirred solution of 2-(4-amino-phenyl)-6-fluoro-2,5-dihydro-pyrazolo[4,3-*c*]quinolin-3-one (Intermediate 2) (30 mg, 0.10 mmol), triethylamine (14 µl, 0.11 mmol) and 4-dimethylaminopyridine (2.4 mg, 0.02 mmol) in dichloromethane (1 ml) at room temperature. The mixture was stirred at room temperature for 16 h. The yellow solid was then filtered and purified by washing sequentially with a saturated solution of sodium bicarbonate (1 ml), ethyl acetate (1 ml) and ethyl alcohol (0.5 ml) and finally dried under suction to give the title compound (10 mg, 26 %) as a bright yellow solid, LCMS m/z 379.36 [M+H]<sup>+</sup> @ R<sub>T</sub> 1.18 min. δ<sub>H</sub>(400 MHz, (CD<sub>3</sub>)<sub>2</sub>SO) 9.89 (1H, s), 8.52 (1H, s), 8.15 (2H, d *J* 9.0 Hz), 8.01 (1H, d *J* 7.0 Hz), 7.69 (2H, d *J* 9.0 Hz) 30 7.57-7.46 (2H, m), 2.46-2.39 (1H, m), 1.69-1.36 (2H, m), 1.11 (3H, d *J* 6.8 Hz), 0.91 (3H, t *J* 7.3 Hz).


The title compound, and compounds of subsequent Examples, were tested in the assay described below in the Assay Section, to determine their activities as inhibitors of the CD80-CD28 interaction. The present  
5 title compound had an activity rating of \*\*\*.

#### Examples 2-49

The following compounds were synthesized by the route described in Example 1, substituting the appropriate acid chloride for (±)-2-methylbutyryl chloride:

#### Example 2

2-Methyl-pentanoic acid [4-(6-fluoro-3-oxo-3,5-dihydro-pyrazolo[4,3-c]quinolin-2-yl)-phenyl]-amide




$\delta_H$  (400 MHz,  $(CD_3)_2SO$ ) 9.92 (1H, s), 8.53 (1H, s), 8.12 (2H, d  $J$  9.2 Hz), 8.05 (1H, d  $J$  7.6 Hz), 7.70 (2H, d  $J$  9.2 Hz), 7.63-7.53 (2H, m), 1.68-1.58 (1H, m), 1.38-1.28 (3H, m), 1.11 (3H, d  $J$  6.6 Hz), 0.91 (3H, t  $J$  7.1 Hz).

Activity \*\*\*

#### Example 3

1-Methyl-1H-pyrrole-2-carboxylic acid [4-(6-fluoro-3-oxo-3,5-dihydro-pyrazolo[4,3-c]quinolin-2-yl)-phenyl]-amide



13


$\delta_H$ (400 MHz,  $(CD_3)_2SO$ ) 9.76 (1H, s), 8.50 (1H, s),  
8.26 (2H, d 9.0 Hz), 7.97-7.94 (1H, m), 7.73 (2H, d  $J$  9.0  
Hz), 7.39-7.28 (2H, m), 7.07-7.01 (2H, m), 3.91 (3H, s).

## 5 Activity \*

Example 4

*N*-[4-(6-Fluoro-3-oxo-3,5-dihydro-pyrazolo[4,3-*c*]quinolin-  
2-yl)-phenyl]-3-methyl-butamide

10



15


$\delta_H$ (400 MHz,  $(CD_3)_2SO$ ) 9.92 (1H, s), 8.52 (1H, s),  
8.14 (2H, d  $J$  9.2 Hz), 8.01 (1H, d  $J$  7.3 Hz), 7.67 (2H,  
20 d  $J$  9.2 Hz), 7.57-7.47 (2H, m), 2.21 (2H, d  $J$  6.8 Hz),  
2.14-2.07 (1H, m), 0.96 (6H, d  $J$  6.6 Hz).

## Activity \*\*

25 Example 5

2-Propyl-pentanoic acid [4-(6-fluoro-3-oxo-3,5-dihydro-  
pyrazolo[4,3-*c*]quinolin-2-yl)-phenyl]-amide

30



35

$\delta_H$ (400 MHz,  $(CD_3)_2SO$ ) 9.93 (1H, s), 8.53 (1H, s),  
8.11 (2H, d  $J$  9.0 Hz), 8.05 (1H, d  $J$  7.8 Hz), 7.70 (2H, d

14

$J$  9.0 Hz), 7.59-7.46 (2H, m), 2.46-2.35 (1H, m), 1.63-1.27 (4H, m), 0.90 (6H, t  $J$  7.1 Hz).


## Activity \*

5

Example 6

5-[4-(6-Fluoro-3-oxo-3,5-dihydro-pyrazolo[4,3-c]quinolin-2-yl) phenylcarbamoyl]-pentanoic acid methyl ester

10



15

$\delta_H$  (400 MHz,  $(CD_3)_2SO$ ) 9.85 (1H, s), 8.47 (1H, s), 8.25 (2H, d  $J$  9.0 Hz), 7.91-7.90 (1H, m), 7.59 (2H, d  $J$  9.0 Hz), 7.29-7.20 (2H, m), 3.61 (3H, s), 2.38-2.28 (4H, m), 1.64-1.50 (4H, m).

20


## Activity \*\*\*

25

Example 7

N-[4-(6-Fluoro-3-oxo-3,5-dihydro-pyrazolo[4,3-c]quinolin-2-yl)-phenyl]-2,2-dimethyl-propionamide

30



35

15


$\delta_H$ (400 MHz,  $(CD_3)_2SO$ ) 9.26 (1H, s), 8.52 (1H, s), 8.15 (2H, d  $J$  9.2 Hz), 8.03 (1H, d  $J$  8.8 Hz), 7.71 (2H, d  $J$  9.2 Hz), 7.56-7.47 (2H, m), 1.26 (9H, s).

5 Activity \*\*

Examples 8 to 28 were also prepared by the method of Example 1 using the appropriate acid chloride:











10

15









Example	X	R	M.S. (MH <sup>+</sup> )	Activity
8	6-F		443.4	**
9	6-F	-CH <sub>2</sub> Cl	371.31	**
10	6-F		389.34	*
11	6-F		485.45	*
12	6-F	CO <sub>2</sub> Me	381.34	**
13	6-F	OE <sub>t</sub>	367.18	
14	6-F		507.43	*
15	6-F		466.41	**
16	6-F	Me	337.36	**
17	6-F	CH(Et)CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> Me	421.46	*
18	6-F	CH(Et) <sub>2</sub>	393.41	***



19	6-F		405.41	**
20	6-F		448.44	**
21	6-F		481.35	**
22	6-F		423.42	***
23	6-F	$(\text{CH}_2)_8\text{CO}_2\text{Me}$	493.51	**
24	6-F	iPr	365.36	***
25	6-F	$\text{CH}_2\text{OCH}_2\text{CH}_2\text{OMe}$	411.4	**
26	6-F	$\text{CH}(\text{Me})(\text{nPr})$	393.42	***
27	6-F	$\text{CH}_2\text{OMe}$	367.24	**
28	6-F		390.33	**
29	6-F	$\text{CH}_2\text{CH}_2\text{CH}_2\text{N}^+(\text{Me})_3$	422.1 (M+)	***
30	6-F	$\text{CH}_2\text{CH}_2\text{CH}_2\text{N}(\text{Me})_2$	408.3	***
31	6-F	$\text{CH}_2\text{NHCH}_2\text{CH}_2\text{CH}_2\text{N}(\text{Me})(\text{Ph})$	499.3	*
32	6-F		485.3	*
33	6-F		505.1	***
34	6-F		517.2	***
35	6-F		477.1	***
36	6-F		457.1	**


17

37	6-F		463.1	**
38	6-F		438.3	**
39	6-F		463.2	***
40	6-F		460.4	**
41	6-F	$\text{CH}_2\text{NHCH}_2\text{CH}_2\text{N}(\text{iPr})_2$	479.4	**
42	6-F		420.2	**
43	H	$\text{CH}(\text{NH}_2)\text{CH}_3$	348.3	**
44	H	$\text{CH}(\text{Me})\text{nPr}$	375.3	*
45	H	iPr	347.3	**
46	6-F	$\text{CH}(\text{NH}_2)\text{CH}_3$	366.3	***
47	H	$\text{CH}(\text{Me})\text{Et}$	361.3	**
48	6-F		529.1	**
49	6-F	$\text{CH}_2\text{N}(\text{Me})\text{CH}_2\text{Ph}$	456.4	**

### Preparation of Intermediate 3

3-(6-Fluoro-3-oxo-3,5-dihydro-pyrazolo[4,3-c]quinolin-2-yl)-benzoic acid

5



10        3-Hydrazinobenzoic acid (1.91 g, 0.013 mol) was added in one portion to a stirred solution of 4-chloro-8-fluoro-quinoline-3-carboxylic acid ethyl ester (2.93 g, 0.011 mol) in n-butanol (60 ml) at room temperature. The solution was heated to reflux for 16 h, cooled to room temperature and the resulting yellow solid filtered, washed with *tert*-butyl methyl ether and then dried. The

15


18

solid was redissolved in a solution of tetrahydrofuran :  
water (2:1; 21 ml) and lithium hydroxide (1.27 g, 0.031  
mol) was then added. After stirring at room temperature  
for 16 h, concentrated hydrochloric acid (3 ml) was added  
5 dropwise to the mixture to precipitate a yellow solid  
which was filtered and dried under vacuum to give the  
title compound (intermediate 3) (2.32 g, 63 %) as a  
bright yellow solid.

10 Preparation of Intermediate 4

3-(6-Fluoro-3-oxo-3,5-dihydro-pyrazolo[4,3-c]quinolin-2-  
yl)-benzoyl chloride

15




Oxalyl chloride (20 ml, 0.2 mol) was added dropwise  
over 2 min to a stirred solution of 3-(6-fluoro-3-oxo-  
20 3,5-dihydro-pyrazolo[4,3-c]quinolin-2-yl)-benzoic acid  
(intermediate 3) (2.0 g, 6.1 mmol) in dichloromethane (10  
ml) at room temperature. *N,N*-Dimethylformamide (50 µl)  
was then added and the resulting mixture heated to 50 °C  
for 1 h. The solution was then cooled to room temperature  
25 and then concentrated in vacuo to leave the title  
compound (intermediate 4) (2.0 g, 96 %) as a beige solid.

Example 50

3-(6-Fluoro-3-oxo-3,5-dihydro-pyrazolo[4,3-c]quinolin-2-  
30 yl)-*N*-(3-methoxy-propyl)-benzamide

35




3-Methoxypropylamine (0.026g, 0.29mmol) was added to a stirred solution of 3-(6-fluoro-3-oxo-3,5-dihydro-pyrazolo[4,3-c]quinolin-2-yl)-benzoyl chloride (intermediate 4) (26 mg 0.29mmol) in tetrahydrofuran (2 ml) and the mixture stirred at room temperature for 15 min. Triethylamine (0.2 ml, 1.4 mmol) was then added and the resulting mixture stirred overnight. 1M Hydrochloric acid (3-4 ml) was added dropwise to precipitate a yellow solid which was filtered and dried under suction to give the amide (79 mg, 0.20 mmol) as a yellow solid, LCMS m/z 395.25 [M+H]<sup>+</sup> @ R<sub>T</sub> 1.04 min;  $\delta_H$ (400 MHz, (CD<sub>3</sub>)<sub>2</sub>SO) 8.59 (1H, m), 8.57 (1H, s), 8.39 (1H, app d *J* 9.3 Hz), 8.08 (1H, app d *J* 7.3 Hz), 7.66-7.53 (5H, m), 3.37-3.33 (4H, m), 3.27 (3H, s), 1.83-1.77 (2H, m).

#### Activity \*\*

##### Example 51

N-Ethyl-3-(6-fluoro-3-oxo-3,5-dihydro-pyrazolo[4,3-c]-quinolin-2-yl)-benzamide



Prepared by the method of Example 53 substituting ethylamine for 3-methoxypropylamine.


$\delta_H$ (400 MHz, (CD<sub>3</sub>)<sub>2</sub>SO) major rotomer quoted; 8.56 (1H, br s), 8.47 (1H, m), 8.21 (2H, d *J* 8.5 Hz), 7.94 (2H, d *J* 8.5 Hz), 3.96 (3H, s), 3.31 (2H, q *J* 7.3 Hz), 2.58 (3H, s), 1.15 (3H, t *J* 7.4 Hz).

#### Activity \*\*

##### Example 52

N-Benzyl-3-(6-fluoro-3-oxo-3,5-dihydro-pyrazolo[4,3-c]-quinolin-2-yl)-benzamide

20



5


Prepared by the method of Example 53 substituting benzylamine for 3-methoxypropylamine.

LCMS m/z 427.16  $[M+H]^+$  @  $R_T$  1.28 min.

10


Activity \*

Examples 53 to 64 were prepared by the method of example 50, using the appropriate amine.



Example	X	R	R'	M.S. (MH <sup>+</sup> )	Activity
53	6-F	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> N(Me) <sub>2</sub>	Me	422.5	*
54	6-F	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> N(Me) <sub>2</sub>	H	408.4	**
55	6-F		H	420.4	*
56	6-F		H	434.4	*
57	6-F		H	448.4	**
58	6-F	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> N(Me) <sub>2</sub>	H	422.4	**
59	6-F	CH <sub>2</sub> CH <sub>2</sub> OMe	H	381.3	**
60	6-F	Et	Et	379.3	*

21

61	6-F	CH <sub>2</sub> CO <sub>2</sub> Me	H	395.2	*
62	6-F	CH <sub>2</sub> CCH	H	361.3	**
63	6-F	CH <sub>2</sub> Ph	Me	427.2	**
64	6-F			463.3	*


Example 65

N-(3-Dimethylamino propyl)-4-(4-cyclopropyl-3-oxo-3,5-dihydro-pyrazolo[4,3-c]quinolin-2-yl)-benzamide

5

Step 1

2-cyclopropyl-4-oxo-1,4-dihydro-quinoline-3-carboxylic acid ethyl ester



10

A solution of 3-cyclopropyl-3-oxo-propionic acid methyl ester (6.2 g, 0.038 mols), 2-amino benzoic acid ethyl ester (4.95 g, 0.03 mols) and *p*-toluene sulfonic acid (0.04 g, 0.2 mmols) in toluene (25 ml) was heated at 125°C for 2h; 15 ml of solvent was then distilled. To the residual orange solution was added sodium ethoxide (2 M, 15 ml) in ethanol (reaction mixture turns red). This red mixture was stirred at 120°C for 2 h; 15 ml of solvent was again distilled. The reaction mixture was left to cool to room temperature, diluted with ethyl acetate (1 litre), extracted with HCl 0.1 M and water. The combined organic extracts were dried over sodium sulfate and concentrated *in vacuo* to leave an orange residue which was washed once with cold ethyl acetate to yield 2-cyclopropyl-4-oxo-1,4-dihydro-quinoline-3-carboxylic acid ethyl ester (3.87 g, 53%) as an off-white solid. LCMS m/z


22

244.14  $[M+H]^+$  @  $R_T$  0.78 min, 89%,  $m/z$  230.11  $[Acid+H]^+$  @  $R_T$  1.27, 11%.

$\delta_H$  (400 MHz,  $(CD_3)_2SO$ ) 11.04 (1 H, s), 8.06 (1 H, dd,  $J_1$  1.1,  $J_2$  8.1), 7.76-7.66 (2 H, m), 7.36 (1 H, td,  $J_1$  1.1,  $J_2$  7.5), 3.89 (3 H, s), 2.16 (1 H, m), 1.18 (4 H, d,  $J$  7.0).

### Step 2


10 4-Chloro-2-cyclopropyl-quinoline-3-carboxylic acid ethyl ester



15 Phosphorus oxychloride (0.77 ml, 0.082 mols) was added in one portion to a suspension of 2-cyclopropyl-4-oxo-1,4-dihydro-quinoline-3-carboxylic acid ethyl ester (1.0 g, 0.041 mols) in acetonitrile and the mixture was heated at 75°C for 90 minutes (becomes a clear solution above  
20 65°C). The resulting light brown solution was poured into saturated sodium bicarbonate (100 ml); the suspension was extracted with ethyl acetate and the combined organic extracts were dried and concentrated *in vacuo* to leave 4-Chloro-2-cyclopropyl-quinoline-3-carboxylic acid ethyl  
25 ester (1.15 g, 106 %) as an off-white solid.  $R_f$  (AcOEt) = 0.73.

### Step 3

30 4-(4-cyclopropyl-3-oxo-3,5-dihydro-pyrazolo[4,3-c]quinolin-2-yl)-benzoic acid



4-Chloro-2-cyclopropyl-quinoline-3-carboxylic acid ethyl ester (1.15 g, 0.0041 mols) and 4-hydrazino-benzoic acid (1.0g, 0.0068 mols) were stirred in ethanol (30 ml) at reflux for 16 h. The bright yellow suspension was diluted with heptane, filtered, washed with cold *t*-butylmethyl ether and left to dry under suction to yield crude solid containing hydrazine. This solid was suspended in 1 M HCl, filtered, washed with water and then dried *in vacuo* to yield 4-(4-cyclopropyl-3-oxo-3,5-dihydro-pyrazolo[4,3-*c*]quinolin-2-yl)-benzoic acid (1.135 g, 80 %) as a yellow solid, LCMS *m/z* 346.20 [M+H]<sup>+</sup> @ *R<sub>T</sub>* 1.05 min: 96% purity.

$\delta_H$  (400 MHz, (CD<sub>3</sub>)<sub>2</sub>SO) 11.4 (1 H, s), 8.43 (2 H, d, *J* 8.1), 8.21 (1 H, dd, *J*<sub>1</sub> 1.2, *J*<sub>2</sub> 8.1), 8.07 (2 H, d, *J* 8.1), 7.92 (1 H, d, *J* 8.1), 7.67 (1 H, t, *J* 6.6), 7.52 (1 H, t, *J* 6.5), 3.43 (1 H, m), 1.59 (2 H, m), 1.43 (2 H, m).


20

#### Step 4

4-(4-cyclopropyl-3-oxo-3,5-dihydro-pyrazolo[4,3-*c*] - quinolin-2-yl)-benzoyl chloride




24



To a suspension of finely ground 4-(4-cyclopropyl-3-oxo-3,5-dihydro-pyrazolo[4,3-c]quinolin-2-yl)-benzoic acid (0.19 g, 0.55 mmol) in dichloromethane (4 ml) was added oxalyl chloride (1.6 ml, 0.01 mol) followed by a drop of dimethyl formamide. The mixture was stirred under nitrogen at 45 °C for 8 h. The solvent was removed in vacuo to yield 4-(4-cyclopropyl-3-oxo-3,5-dihydro-pyrazolo[4,3-c]quinolin-2-yl)-benzoyl chloride as a pale yellow solid, LCMS m/z [M+MeOH-Cl]<sup>+</sup> @ R<sub>T</sub> 1.46 min: 95% purity. Used without further purification.

#### Step 5

N-(3-Dimethylamino propyl)-4-(4-cyclopropyl-3-oxo-3,5-dihydro-pyrazolo[4,3-c]quinolin-2-yl)-benzamide



To a partial solution of 4-(4-cyclopropyl-3-oxo-3,5-dihydro-pyrazolo[4,3-c]quinolin-2-yl)-benzoyl chloride (0.1 g, 0.28 mmol) in tetrahydrofuran (6 ml) under nitrogen was added a solution of 3-dimethylamino-propyl

25


amine (0.03 g, 0.3 mmol) in tetrahydrofurane (3 ml). The mixture was stirred at RT for 3 h. The solvent was removed under reduced pressure and the yellow solid was washed with a little saturated sodium bicarbonate, water and dried under vacuo to yield *N*-(3-Dimethylamino propyl)-4-(4-cyclopropyl-3-oxo-3,5-dihydro-pyrazolo[4,3-*c*]-quinolin-2-yl)-benzamide (57 mg, 47 %) as a yellow solid. LCMS *m/z* 430.11 [M+H]<sup>+</sup> @ *R<sub>T</sub>* 0.99 min: 100% purity.

10           Activity \*\*\*

Preparation of Intermediate 5

4-(6-Fluoro-3-oxo-3,5-dihydro-pyrazolo[4,3-*c*]quinolin-2-yl)-benzoyl chloride

15




To a suspension of finely ground 4-(6-Fluoro-3-oxo-3,5-dihydro-pyrazolo[4,3-*c*]quinolin-2-yl)-benzoic acid ( 1.1 g. 3.4 mmol) in dichloromethane (6 ml) was added oxalyl chloride (2.4 ml, 29 mmol) followed by a drop of dimethyl formamide. The mixture was stirred under nitrogen at 45 °C for 3 h. The solvent was removed in vacuum to yield 4-(6-Fluoro-3-oxo-3,5-dihydro-pyrazolo[4,3-*c*]quinolin-2-yl)-benzoyl chloride (1.15 g, quantitative) as a pale yellow solid that was used without further purification.

Example 66

N-(3-Dimethylamino propyl)-4-(6-fluoro-3-oxo-3,5-dihydro-pyrazolo[4,3-c]quinolin-2-yl)-benzamide hydrochloride

5




To a partial solution of 4-(6-Fluoro-3-oxo-3,5-dihydro-pyrazolo[4,3-c]quinolin-2-yl)-benzoyl chloride (0.1 g,  
10 0.3 mmol) in tetrahydrofuran (5 ml) under nitrogen was added a solution of 3-dimethylamino-propyl amine (0.03 g, 0.3 mmol) in tetrahydrofuran. The mixture was stirred at rt for 90 minutes. The solvent was removed under reduced pressure and the yellow solid was purified via  
15 FCC silica gel (gradient elution, MeOH:H<sub>2</sub>O, Fluka C<sub>18</sub> reverse phase) to yield N-(3-Dimethylamino propyl)-4-(6-fluoro-3-oxo-3,5-dihydro-pyrazolo[4,3-c]quinolin-2-yl)-benzamide hydrochloride (70 mg, 53 %) as a yellow solid.













20 LCMS m/z 408.39 [M+H]<sup>+</sup> @ R<sub>T</sub> 0.89 min: 90% purity.

Activity \*\*\*












Exmaples 67 - 141 were prepared analogously from the  
25 appropriate benzoyl chloride and the appropriate amine


















Example	X	Z	W	R	R'	M.S. (MH <sup>+</sup> )	Activ ity
67	6-F	H	H	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -		391.3	**
68	6-F	H	H	-CH <sub>2</sub> Phenyl	H	413.2	***
69	6-F	H	H	-CH <sub>2</sub> Phenyl	Me	427.3	**
70	6-F	H	H	-CH <sub>2</sub> CH <sub>2</sub> OMe	H	381.2	***
71	6-F	H	H	-CH <sub>2</sub> CH <sub>2</sub> N(Me) <sub>2</sub>	H	394.3	***
72	6-F	H	H	-CH <sub>2</sub> CO <sub>2</sub> Me	H	395.3	***
73	6-F	H	H	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> OMe	H	395.2	***
74	6-F	H	H	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> N(Me) <sub>2</sub>	H	408.3	***
75	6-F	H	H		H	431.3	**
76	6-F	H	H		H	419.2	**
77	6-F	H	H	Et	H	351.2	***
78	6-F	H	H	Et	Et	379.3	**
79	6-F	H	H		H	420.4	***
80	6-F	H	H	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> N(Me) <sub>2</sub>	Me	422.4	***
81	6-F	H	H	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> N(Me) <sub>2</sub>	H	422.4	***
82	6-F	H	H		H	448.5	***
83	6-F	H	H		H	434.4	***
84	6-F	H	H		H	525.3	***
85	6-F	H	H	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> N(Me) <sub>2</sub>	H	450.3	***
86	H	H	H	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> N(Me) <sub>2</sub>	H	390.2	***
87	H	H	H	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> N(Me) <sub>2</sub>	H	432.1	**



















88	H	H	H	$-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{N}(\text{Et})_2$	H	432.2	**
89	H	H	H	$-\text{CH}_2\text{CH}_2\text{CH}_2\text{N}(\text{Me})_2$	Me	404.2	**
90	6-F	H	2-Cl	$-\text{CH}_2\text{CH}_2\text{CH}_2\text{N}(\text{Me})_2$	H	442.1	**
91	H	H	H		H	416.1	**
92	H	H	H		H	573.0	**
93	H	H	H		H	445.1	**
94	H	H	H		H	507.1	**
95	6-F	H	H		H	591.0	***
96	H		H	$-\text{CH}_2\text{CH}_2\text{CH}_2\text{N}(\text{Me})_2$	H	430.1	***
97	6-F	H	H		H	464.1	***
98	6-F	H	H		H	463.1	***
99	6-F	H	3-Cl		H	482.1	**
100	6-F	H	2-Cl		H	497.1	**
102	6-F	H	2-Cl	$-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{N}(\text{Et})_2$	H	484.1	**
103	6-F	H	3-Cl	$-\text{CH}_2\text{CH}_2\text{CH}_2\text{N}(\text{Me})_2$	H	442.1	**
104	H		H		H	470.4	***

29

105	6-F	H	H			516.3	*
106	6-F	H	H		H	470.3	***
107	6-F	H	H	$-\text{CH}_2\text{CH}_2\text{N}(\text{iPr})_2$	H	451.4	***
108	6-F	H	2-Cl		H	496.2	**
109	6-F	H	H		H	456.1	***
110	6-F	H	2-Cl	$-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{N}(\text{Me})_2$	H	456.1	**
111	6-F	H	H			406.2	**
112	6-F	H	H		H	462.1	***
113	6-F	H	H		H	436.1	***
114	6-F	H	H		H	434.4	***
115	6-F	H	H		H	476.1	***
116	6-F	H	H		H	496.1	***
117	6-F	H	H		H	436.3	***

30

118	6-F	H	H		H	462.3	***
119	6-F	H	H		H	428.1	**
120	6-F	H	H	-CH <sub>2</sub> CH <sub>2</sub> SEt	H	411.3	***
121	6-F	H	H		H	448.3	**
122	6-F	H	H		H	431.3	***
123	6-F	H	H		H	434.3	**
124	6-F	H	H	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> N(Et) <sub>2</sub>	H	450.4	***
125	6-F		H		H	536.1	***
126	6-F		H		H	516.2	***
127	6-F	H	H		H	428.3	*
128	6-F	H	H	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> SMe	H	411.3	**
129	H		H		H	498.5	***
130	6-F		H		H	488.4	***
131	6-F	H	H		H	446.3	***


132	6-F		H	$-\text{CH}_2\text{CH}_2\text{CH}_2\text{N}(\text{Me})_2$	H	448.2	***
133	6-F		H		H	502.3	***
134	6-F		H		H	486.3	***
135	6-F		H	$-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{N}(\text{Et})_2$	H	490.3	***
136	6-F		H		H	546.2	**
137	6-F		H		H	631.2	***
138	6-F		H		H	468.2	**
139	6-F		H		H	468.2	*
140	6-F		H		H	476.2	***
141	6-F		H		H	474.3	***



Example 142

{3-[4-(6-Fluoro-3-oxo-3,5-dihydro-pyrazolo[4,3-c]quinolin-2-yl)-phenyl]-ureido} acetic acid ethyl ester

5




10 Ethyl cyanatoacetate (31 mg, 0.24 mmol) was added in one portion to a stirred solution of 2-(4-aminophenyl)-6-fluoro-2,5-dihydropyrazolo[4,3-c]quinolin-3-one (intermediate 2) (50 mg, 0.17 mmol) in *N,N*-dimethylformamide (2 ml) and the mixture stirred at room temperature for 16  
15 h. Water (1 ml) was then added to the mixture to precipitate a solid, which was filtered, washed with water (1 ml) and then ethyl acetate (1 ml) and finally dried by suction to leave the urea as a yellow solid, LCMS *m/z* 424.40  $[M+H]^+$  @ *R<sub>T</sub>* 1.06 min.

20

Activity \*\*\*


Examples 143 and 144

25



Example 143  
LCMS *m/z* 438.41  $[M+H]^+$  @ RT 1.13 min.

Activity \*\*




Example 144  
LCMS *m/z* 514.46  $[M+H]^+$  @ RT 1.35 min.



Activity \*

The following compounds were synthesised by the method of Example 142, substituting the appropriate isocyanate, isothiocyanate or chloroformate for ethyl cyanatoacetate.


5



Example	X	Z	Y	R	A	M.S. (MH <sup>+</sup> )	Activ ity
144	6-F	H	O	iPr	NH	380.3	***
145	6-F	H	O	nPr	NH	380.3	***
146	6-F	H	O	tBu	NH	394.4	***
147	6-F	H	O	Ph	NH	414.3	**
148	6-F	H	S		NH	394.3	**
149	6-F	H	S		NH	436.4	*
150	6-F	H	O	tBu	O	395.3	***
151	6-F	H	O	Et	O	367.2	**
152	6-F	H	O	CH <sub>2</sub> CH <sub>2</sub> N (Me) <sub>2</sub>	O	410.2	***
153	H		O	Me	O	375.3	**
154	6-F	H	O	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> N (Me) <sub>2</sub>	O	424.1	***
155	6-F	H	O		O	512.3	**
156	6-F	H	S	nPentyl	NH	424.4	**
157	6-F	H	S	CH (CH <sub>3</sub> ) CH (CH <sub>3</sub> ) CH <sub>3</sub>	NH	424.4	**
158	6-F	H	O	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> N (Et) <sub>2</sub>	NH	465.4	***
159	H	H	O	nPr	NH	362.3	***
160	H	H	S		NH	376.1	**

161	6-F	H	O	$\text{CH}_2\text{CH}_2\text{CH}_2\text{N}(\text{Me})_2$	NH	423.3	***
162	H	H	O		NH	434.5	***
163	6-F	H	O	$\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{N}(\text{Me})_2$	NH	437.2	***
164	6-F	H	O		NH	463.5	***

Intermediate 6: Preparation of methyl 4-oxothiochromane-  
5 3-carboxylate




Dry tetrahydrofuran (60 ml) was cooled under nitrogen  
10 atmosphere to  $-50$  to  $-60^\circ\text{C}$ . 1M Lithium  
bis(trimethylsilyl)amide solution in hexane (56 ml, 56  
mmol) was added. The temperature was kept at  $-50$  to  $-60^\circ\text{C}$   
and thiochroman-4-one was added dropwise over 20 min.  
Stirring was continued at low temperature for 60 min.  
15 Methyl cyanoformate (4.84 ml, 60.9 mmol) was added  
dropwise over 5 min to the reaction mixture. The obtained  
suspension was stirred at  $-50$  to  $-60^\circ\text{C}$  for 80 min and  
then allowed to warm up to room temperature. Saturated  
ammonium chloride solution (100 ml) was added. The phases  
20 were separated, the aqueous phase extracted with ethyl  
acetate (2 x 100 ml). The combined organic phases were  
washed with water (50 ml), dried over magnesium sulphate,  
filtered and concentrated under vacuum. An orange oil was  
obtained and purified by column chromatography. The title

35

compound was isolated as a yellow solid (4.70 g, 21.1 mmol, 42%). LCMS: m/z 221 [M-H]<sup>+</sup>.


5 Intermediate 7: Preparation of 4-(3-Oxo-3a,4-dihydro-3H-thiochromeno[4,3-c]pyrazol-2-yl)-benzoic acid



10 4-Oxothiochromane-3-carboxylate (0.50 g, 2.25 mmol) and hydrazinobenzoic acid (0.377 g, 2.48 mmol) were mixed in acetic acid (6 ml). The mixture was heated to reflux for 30 min. Excess acetic acid was distilled off to give a brown oil. Diethylether was added, a precipitate formed  
15 which was collected by filtration and dried under vacuum. The crude product was isolated as a red/brown solid (797 mg). LCMS: m/z 325 [M+H]<sup>+</sup>. No purification was carried out.

20

Intermediate 8: Preparation of 4-(3-oxothiochromeno[4,3-c]pyrazol-2(3H)-yl)benzoic acid




Crude 4-(3-Oxo-3a,4-dihydro-3H-thiochromeno[4,3-c]pyrazol-2-yl)-benzoic acid (250 mg, 0.77 mmol) was dissolved in dimethyl sulphoxide (6 ml). O-Chloranil (189 mg, 0.77 mmol) was added and the mixture was stirred at room temperature overnight. Water (20 ml) was added and the solids were collected by filtration and washed with water. The filter cake was triturated with toluene, filtered and dried under vacuum. The title compound was isolated as a dark brown solid (230 mg, 0.71 mmol, 92%). LCMS: m/z 323 [M+H]<sup>+</sup>

Alternatively crude 4-(3-Oxo-3a,4-dihydro-3H-thiochromeno[4,3-c]pyrazol-2-yl)-benzoic acid can be stirred in dimethyl sulphoxide under exposure to air. It was found that air oxidation provides clean product, however the reaction is much slower.

20

#### Example 165

Preparation of *N*-[3-(dimethylamino)propyl]-4-(3-oxothiochromeno[4,3-c]pyrazol-2(3H)-yl)benzamide




4-(3-oxothiochromeno[4,3-c]pyrazol-2(3H)-yl)benzoic acid (55 mg, 0.17 mmol) was suspended in anhydrous dimethyl acetamide (1 ml). Diisopropyl-ethyl amine (46.5 mg, 0.36 mmol, 62  $\mu$ l) was added followed by 3-dimethylaminopropylamine (17.5 mg, 0.17 mmol) and [(benzotriazol-1-yloxy)-dimethylamino-methylene]-dimethyl-ammonium hexafluoro phosphate (65 mg, 0.17 mmol). The mixture was stirred at room temperature for 4 h and was purified by preparative HPLC. The title compound was isolated as a brown solid. LCMS: m/z 407 [M+H]<sup>+</sup>

15 Activity \*\*

#### Example 166

Preparation of N-[(cyclohexylamino)propyl]-4-(3-oxothiochromeno[4,3-c]pyrazol-2(3H)-yl)benzamide



38


The reaction was carried out as described above. LCMS:  
m/z 461 [M+H]<sup>+</sup>

Activity \*\*\*

5

Example 167

Preparation of N-(pyrrolidin-1-yl-butyl)-4-(3-oxothiochromeno[4,3-c]pyrazol-2(3H)-yl)benzamide



10

The reaction was carried out as described above. LCMS:  
m/z 447 [M+H]<sup>+</sup>


15 Activity \*

Example 168

Preparation of 4-(3-oxothiochromeno[4,3-c]pyrazol-2(3H)-yl)-N-1,2,2,6,6-pentamethylpiperidin-4-ylbenzamide

20

39




The reaction was carried out as described above. LCMS:  
m/z 475 [M+H]<sup>+</sup>

5

Activity \*\*

Intermediate 9: Preparation of 3-[(2-fluorophenyl)sulfanyl]propanoic acid

10



2-Fluorothiophenol (5.0 g, 39 mmol) was dissolved in  
15 tetrahydrofuran (50 ml) under a nitrogen atmosphere.  
Triethylamine (3.94 g, 5.33 ml, 85.8 mmol) was added.  
Acrylic acid (2.81 g, 2.67 ml, 39 mmol) was dissolved in  
tetrahydrofuran and added dropwise to the reaction  
solution over 2 h at room temperature. The mixture was  
20 stirred at room temperature overnight. 1M Hydrochloric  
acid (50 ml) was added and the phases were separated. The  
aqueous phase was washed with ethyl acetate (2 x 50 ml).  
The combined organic phases were dried over magnesium  
sulphate, filtered and concentrated under vacuum. A




40

yellow oil was obtained which solidified upon storage at room temperature. The solid was triturated with hexane, filtered and dried under vacuum. The title compound was isolated as an off-white solid (4.19 g, 20.9 mmol, 54%).

5

Intermediate 10: Preparation of 8-fluoro-2,3-dihydro-4H-thiochromen-4-one




10

3-[(2-Fluorophenyl)sulfanyl]propanoic acid (4.0 g, 20 mmol) was mixed with concentrated sulphuric acid (20 ml) at 0-5°C. The reaction solution was stirred at 0 to 5°C for 3 h then allowed to warm up to room temperature overnight. The mixture was quenched dropwise into ice to give a white suspension. The aqueous phase was extracted with ethyl acetate (1 x 200 ml, 1 x 100 ml). The combined organic phases were washed with saturated sodium bicarbonate solution (1 x 50 ml), water (1 x 50 ml), 1M hydrochloric acid (50 ml) and water (2 x 50 ml). The organic phase was dried over magnesium sulphate, filtered and concentrated under vacuum. The title compound was isolated as a yellow solid (2.10 g, 11.5 mmol, 58%).

25


Intermediate 11: Preparation of methyl 8-fluoro-4-oxothiochromane-3-carboxylate



1M Lithium hexamethyldisilazide solution in hexane (13.2 ml) was dissolved in anhydrous tetrahydrofuran (20 ml) under nitrogen atmosphere. The solution was cooled to -78°C. 8-Fluoro-2,3-dihydro-4H-thiochromen-4-one (2.00 g, 11 mmol) was dissolved in tetrahydrofuran (40 ml), the solution was transferred to the dropping funnel and added dropwise over 30 min to the reaction mixture maintaining the temperature below -60°C. An orange clear solution was obtained which was stirred at -78°C to -65°C for 2 h. Methyl cyanoformate (0.935 g, 0.87 ml) was dissolved in tetrahydrofuran (2 ml) and added dropwise to the reaction solution. Stirring was continued at low temperature for 1 h, the mixture was then allowed to warm to room temperature. Saturated ammonium chloride solution (20 ml) and water (10 ml) were added, the phases mixed for 5 min and separated. The aqueous phase was washed with ethyl acetate (2 x 100 ml) and the combined organic phases were dried over magnesium sulphate. The mixture was filtered and the solvent removed under vacuum to give an orange oil. The crude oil was purified by column chromatography; mobile phase: hexanes, gradient to hexanes / ethyl acetate [90:10]. The title compound was isolated as a yellow solid (1.19 g, 4.95 mmol, 45%).

Intermediate 12: Preparation of 4-(6-fluoro-3-oxothiochromeno[4,3-c]pyrazol-2(3H)-yl)benzoic acid

42




Methyl 8-fluoro-4-oxothiochromane-3-carboxylate (1.19 g, 4.95 mmol) and 4-hydrazinobenzoic acid (755 mg, 4.95 mmol) were mixed with glacial acetic acid (10 ml). The mixture was heated to reflux for 4 h. Excess acetic acid was removed under vacuum to give an orange oil. Ethyl acetate (10 ml) was added and the mixture sonicated. Precipitation of an orange solid was observed. The solids were collected by filtration and washed with ethyl acetate. The filter cake was taken up in dimethyl sulfoxide (10 ml) and air-oxidised at room temperature for one week. Water (20 ml) was added to the reaction mixture, the solids were collected by filtration, slurried in ethyl acetate, filtered and dried under vacuum. The title compound was isolated as an orange powder (175 mg, 0.51 mmol, 10%). LCMS: m/z 341.

#### 20 Example 169

Preparation of *N*-[3-(dimethylamino)propyl]-4-(6-fluoro-3-oxothiochromeno[4,3-*c*]pyrazol-2(3*H*)-yl)benzamide

43




4-(6-Fluoro-3-oxothiochromeno[4,3-c]pyrazol-2(3H)-yl)benzoic acid (41 mg, 0.12 mmol) was dissolved in anhydrous dimethyl-acetamide (1 ml). Diisopropyl-ethyl amine (46 mg, 0.36 mmol, 62  $\mu$ l) was added followed by [(benzotriazol-1-yloxy)-dimethylamino-methylene]-dimethyl-ammonium hexafluoro phosphate (65 mg, 0.17 mmol) and 3-dimethylaminopropylamine (12 mg, 0.12 mmol). The mixture was stirred at room temperature overnight and purified by preparative HPLC. The title compound was isolated as a brown solid. LCMS: m/z 425 [M+H]<sup>+</sup>.

Activity \*\*

15

#### Example 170

Preparation of N-[(cyclohexylamino)propyl]-4-(6-fluoro-3-oxothiochromeno[4,3-c]pyrazol-2(3H)-yl)benzamide




The reaction was carried out as described above. LCMS:  
m/z 479 [M+H]<sup>+</sup>.

5 Activity \*\*

Example 171

Preparation of *N*-(pyrrolidin-1-yl-butyl)-4-(6-fluoro-3-oxothiochromeno[4,3-*c*]pyrazol-2(3*H*)-yl)benzamide

10



The reaction was carried out as described above. LCMS:  
m/z 465 [M+H]<sup>+</sup>.


15

Activity \*\*\*

Example 173

Preparation of 4-(6-fluoro-3-oxothiochromeno[4,3-*c*]pyrazol-2(3*H*)-yl)-*N*-1,2,2,6,6-pentamethylpiperidin-4-ylbenzamide

20



The reaction was carried out as described above. LCMS:  
m/z 493 [M+H]<sup>+</sup>

5

Activity \*\*\*

#### Assay Section

10        The examples described above were tested in a cell free Homogenous Time Resolved Fluorescence (HTRF) assay to determine their activity as inhibitors of the CD80-CD28 interaction.

15        In the assay, europium and allophycocyanin (APC) are associated with CD28 and CD80 indirectly (through antibody linkers) to form a complex, which brings the europium and APC into close proximity to generate a signal. The complex comprises the following six proteins: fluorescent label 1, linker antibody 1, CD28 fusion protein, 20 CD80 fusion protein, linker antibody 2, and fluorescent label 2. The table below describes these reagents in greater detail.

Fluorescent label 1	Anti-Rabbit IgG labelled with Europium (1µg/ml)
Linker antibody 1	Rabbit IgG specific for mouse Fc fragment (3µg/ml)
CD28 fusion protein	CD28 - mouse Fc fragment fusion protein (0.48µg/ml)

CD80 fusion protein	CD80 mouse Fab fragment (C215) fusion protein (1.9 $\mu$ g/ml)
Linker antibody 2	G $\alpha$ M $\kappa$ -biotin: biotinylated goat IgG specific for mouse kappa chain (2 $\mu$ g/ml)
Fluorescent label 2	SA-APC: streptavidin labelled allophycocyanin (8 $\mu$ g/ml)

On formation of the complex, europium and APC are brought into proximity and a signal is generated.

Non-specific interaction was measured by substituting a mouse Fab fragment (C215) for the CD80 mouse Fab fragment fusion protein (1.9 $\mu$ g/ml). The assay was carried out in black 384 well plates in a final volume of 30 $\mu$ l. Assay buffer: 50mM Tris-HCl, 150mM NaCl pH7.8, containing 0.1% BSA (w/v) added just prior to use.

Compounds were added to the above reagents in a concentration series ranging between 100 $\mu$ M - 1.7nM. The reaction was incubated for 4 hours at room temperature. Dual measurements were made using a Wallac Victor 1420 Multilabel Counter. First measurement: excitation 340nm, emission 665nm, delay 50 $\mu$ s, window time 200 $\mu$ s. second measurement: excitation 340nm, emission 615nm, delay 50 $\mu$ s, window time 200 $\mu$ s. Counts were automatically corrected for fluorescence crossover, quenching and background.

By way of illustration, the EC<sub>50</sub> results for the compounds of Examples 15, 21, 29, 35 and 83 were 8  $\mu$ M, 1.9  $\mu$ M, 950 nM, 148nM and 90nM respectively. For convenience, the EC<sub>50</sub> activities of compounds tested are recorded above in summary form as:


EC<sub>50</sub>: \* = >10  $\mu$ M, \*\* = 1-10  $\mu$ M, \*\*\* = <1  $\mu$ M.

## CLAIMS

1. A compound of formula (I) or a pharmaceutically or veterinarily acceptable salt thereof:

5

10



wherein

R<sub>1</sub> and R<sub>3</sub> independently represent H; F; Cl; Br; -NO<sub>2</sub>; -CN; C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted by F or Cl; or C<sub>1</sub>-C<sub>6</sub> alkoxy optionally substituted by F;

R<sub>2</sub> represents H, or optionally substituted C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>3</sub>-C<sub>7</sub> cycloalkyl or optionally substituted phenyl;

Y represents -O-, -S-, N-oxide, or -N(R<sub>5</sub>)- wherein R<sub>5</sub> represents H or C<sub>1</sub>-C<sub>6</sub> alkyl;

X represents a bond or a divalent C<sub>1</sub>-C<sub>6</sub> alkylene radical;

R<sub>4</sub> represents -C(=O)NR<sub>6</sub>R<sub>7</sub>, -NR<sub>7</sub>C(=O)R<sub>6</sub>, -NR<sub>7</sub>C(=O)OR<sub>6</sub>, -NHC(=O)NHR<sub>6</sub> or -NHC(=S)NHR<sub>6</sub> wherein

R<sub>6</sub> represents H, or a radical of formula -(Alk)<sub>b</sub>-Q wherein b is 0 or 1 and

Alk is an optionally substituted divalent straight chain or branched C<sub>1</sub>-C<sub>12</sub> alkylene, C<sub>2</sub>-C<sub>12</sub> alkenylene or C<sub>2</sub>-C<sub>12</sub> alkynylene radical which may be interrupted by one or more non-adjacent -O-, -S- or -N(R<sub>8</sub>)- radicals wherein R<sub>8</sub> represents H or C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>3</sub>-C<sub>4</sub> alkenyl, C<sub>3</sub>-C<sub>4</sub> alkynyl, or C<sub>3</sub>-C<sub>6</sub> cycloalkyl, and

Q represents H; -CF<sub>3</sub>; -OH; -SH; -NR<sub>8</sub>R<sub>8</sub> wherein each R<sub>8</sub> may be the same or different; an ester group; or an optionally substituted phenyl, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, C<sub>5</sub>-C<sub>7</sub> cycloalkenyl or heterocyclic ring having from 5 to 8 ring atoms; and



R<sub>7</sub> represents H or C<sub>1</sub>-C<sub>6</sub> alkyl; or when taken together with the atom or atoms to which they are attached R<sub>6</sub> and R<sub>7</sub> form an optionally substituted heterocyclic ring having from 5 to 8 ring atoms.

5        2. A compound as claimed in claim 1 wherein R<sub>1</sub> is H, F, Cl, methyl or methoxy.

3. A compound as claimed in claim 1 or claim 2 wherein R<sub>2</sub> is H, methyl, methoxy, cyclopropyl, phenyl, or fluoro-, chloro-, methyl, or methoxy-substituted phenyl.

10       4. A compound as claimed in any of the preceding claims wherein R<sub>3</sub> is H, F, Cl, methyl, methoxy, or methylenedioxy.

5. A compound as claimed in any of the preceding claims wherein Y is -O-, -S-, or -N(R<sub>5</sub>)- wherein R<sub>5</sub>  
15 represents H or methyl.

6. A compound as claimed in any of the preceding claims wherein X is a bond, or a -CH<sub>2</sub>- or -CH<sub>2</sub>CH<sub>2</sub>- radical.

7. A compound as claimed in any of the preceding  
20 claims wherein R<sub>4</sub> represents -C(=O)NHR<sub>6</sub>, -NR<sub>7</sub>C(=O)R<sub>6</sub>, -NR<sub>7</sub>C(=O)OR<sub>6</sub>, -NHC(=O)NHR<sub>6</sub> or -NHC(=S)NHR<sub>6</sub> and in these R<sub>6</sub> is H or a radical of formula -Alk<sub>b</sub>-Q wherein

b is 0 or 1 and

Alk is a -(CH<sub>2</sub>)<sub>n</sub>-, -CH((CH<sub>2</sub>)<sub>m</sub>CH<sub>3</sub>)(CH<sub>2</sub>)<sub>n</sub>-,  
25 -CH((CH<sub>2</sub>)<sub>m</sub>CH<sub>3</sub>)((CH<sub>2</sub>)<sub>p</sub>CH<sub>3</sub>)(CH<sub>2</sub>)<sub>n</sub>-, -(CH<sub>2</sub>)<sub>n</sub>-O-(CH<sub>2</sub>)<sub>m</sub>-,  
or -(CH<sub>2</sub>)<sub>n</sub>-O-(CH<sub>2</sub>)<sub>n</sub>-O-(CH<sub>2</sub>)<sub>m</sub>-, radical where n is 1, 2, 3  
or 4 and m and p are independently 0, 1, 2, 3 or 4, and  
Q represents H, -OH, -COOCH<sub>3</sub> phenyl, cyclopropyl,  
cyclopentyl, cyclohexyl, pyridyl, furyl, thienyl, or  
30 oxazolyl. and

R<sub>7</sub> is H, or when taken together with the nitrogen atom to which they are attached R<sub>6</sub> and R<sub>7</sub> form a pyrrolidine-2-one or pyrrolidine-2,5-dione ring.

8. A compound as claimed in claim 1 wherein R<sub>1</sub> is  
35 H, F, or Cl; R<sub>2</sub> is H; R<sub>3</sub> is H, F, or Cl; Y is -NH-; X is a bond; and R<sub>4</sub> represents -C(=O)NHR<sub>6</sub>, -NR<sub>7</sub>C(=O)R<sub>6</sub>, -NR<sub>7</sub>C(=O)OR<sub>6</sub> or -NHC(=O)NHR<sub>6</sub> wherein:

$R_6$  is H or a radical of formula  $-Alk_b-Q$  wherein  $b$  is 0 or 1 and

$Alk$  is a  $-(CH_2)_n-$ ,  $-CH((CH_2)_mCH_3)(CH_2)_n-$ ,  
5  $-CH((CH_2)_mCH_3)((CH_2)_pCH_3)(CH_2)_n-$ ,  $-(CH_2)_n-O-(CH_2)_m-$ ,  
or  $-(CH_2)_n-O-(CH_2)_n-O-(CH_2)_m-$ , radical where  $n$  is 1, 2, 3  
or 4 and  $m$  and  $p$  are independently 0, 1, 2, 3 or 4, and  
 $Q$  represents H,  $-OH$ ,  $-COOCH_3$  phenyl, cyclopropyl,  
cyclopentyl, cyclohexyl, pyridyl, furyl, thienyl, or  
oxazolyl. and

10  $R_7$  is H, or when taken together with the nitrogen  
atom to which they are attached  $R_6$  and  $R_7$  form a pyrro-  
lidine-2-one or pyrrolidine-2,5-dione ring.

9. A compound as claimed in claim 1 wherein  $R_1$  is  
F,  $R_2$  is H or cyclopropyl,  $R_3$  is H,  $X$  is a bond, and  $R_4$  is  
15  $-C(=O)NHR_6$ ,  $-NRHC(=O)R_6$ , or  $-NHC(=O)NHR_6$ .

10. N-(3-Dimethylamino propyl)-4-(4-cyclopropyl-3-  
oxo-3,5-dihydro-pyrazolo[4,3-c]quinolin-2-yl)-benzamide,  
or pharmaceutically or veterinarily acceptable salt  
thereof.

20 11. A compound as claimed in any of claims 1 to 10  
for use in the treatment of conditions which benefit from  
immunomodulation.

12. The use of a compound as claimed in any of  
claims 1 to 10 in the manufacture of a medicament for the  
25 treatment of conditions which benefit from immunomodu-  
lation.

13. A method of immunomodulation in humans and non-  
human primates, comprising administration to a subject in  
need of such treatment an immunomodulatory effective dose  
30 of a compound as claimed in any of claims 1 to 10.

14. A pharmaceutical or veterinary composition com-  
prising a compound as claimed in any of claims 1 to 10  
together with a pharmaceutically or veterinarily accep-  
table excipient or carrier.

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 2003/001805

## A. CLASSIFICATION OF SUBJECT MATTER

IPC7: C07D 471/06, A61K 31/437, A61P 37/02

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7: C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CHEM. ABS DATA, WPI DATA, EPO-INTERNAL

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
PX	WO 03004495 A1 (ACTIVE BIOTECH AB), 16 January 2003 (16.01.2003)  --	1-14
X	WO 9111448 A1 (THE BOOTS COMPANY PLC), 8 August 1991 (08.08.1991), see the claims  --	1-14
X	WO 9734893 A1 (ASTRA PHARMACEUTICALS LTD.), 25 Sept 1997 (25.09.1997)  -- -----	1-14



Further documents are listed in the continuation of Box C.



See patent family annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

5 February 2004

Date of mailing of the international search report

17 -02- 2004

Name and mailing address of the ISA/

Swedish Patent Office

Box 5055, S-102 42 STOCKHOLM

Facsimile No. +46 8 666 02 86

Authorized officer

Carolina Gómez Lagerlöf/EÖ

Telephone No. +46 8 782 25 00

# INTERNATIONAL SEARCH REPORT

International application No.  
**PCT/SE 2003/001805**

## Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: **13**  
because they relate to subject matter not required to be searched by this Authority, namely:  
**see next sheet**
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 2003/001805

Box No. IV Text of the abstract (Continuation of item 5 of the first sheet)

Claim 13 relates to a method of treatment of the human or animal body by surgery or by therapy/a diagnostic method practised on the human or animal body/Rule 39.1(iv). Nevertheless, a search has been executed for this claim. The search has been based on the alleged effects of the compound/composition.

## INTERNATIONAL SEARCH REPORT

Information on patent family members

24/12/2003

International application No.

PCT/SE 2003/001805

WO	03004495	A1	16/01/2003	AU	1290302	A	15/05/2002
				SE	0102404	D	00/00/0000
				US	6642249	B	04/11/2003
				US	2003022913	A	30/01/2003

---

WO	9111448	A1	08/08/1991	AU	7220991	A	21/08/1991
				BR	9105984	A	10/11/1992
				CA	2074841	A	03/08/1991
				CS	9100238	A	15/09/1991
				EP	0539372	A	05/05/1993
				FI	923486	A	31/07/1992
				GB	9002314	D	00/00/0000
				HU	9202510	D	00/00/0000
				IE	910365	A	14/08/1991
				IL	97114	D	00/00/0000
				IN	171735	A	26/12/1992
				JP	5505180	T	05/08/1993
				MX	24401	A	01/10/1993
				NO	923037	A	30/09/1992
				PT	96656	A	31/10/1991
				ZA	9100764	A	30/10/1991
				GB	9002315	D	00/00/0000
				GB	9002425	D	00/00/0000

---

## INTERNATIONAL SEARCH REPORT

Information on patent family members

24/12/2003

International application No.

PCT/SE 2003/001805

WO	9734893	A1	25/09/1997	AU	712141	B	28/10/1999
				AU	2186797	A	10/10/1997
				BR	9708103	A	27/07/1999
				CA	2247814	A	25/09/1997
				CN	1218472	A	02/06/1999
				CZ	9802977	A	17/03/1999
				EE	9800298	A	15/02/1999
				EP	0888347	A	07/01/1999
				GB	9605803	D	00/00/0000
				ID	16283	A	00/00/0000
				IL	126271	D	00/00/0000
				JP	2000506884	T	06/06/2000
				NO	984290	A	27/10/1998
				NZ	331614	A	28/07/2000
				PL	328921	A	01/03/1999
				SK	118798	A	10/03/1999
				TR	9801861	T	00/00/0000
				ZA	9702150	A	22/09/1997
				GB	9610474	D	00/00/0000
				GB	9610894	D	00/00/0000
				AU	719553	B	11/05/2000
				AU	2170397	A	22/10/1997
				BR	9708478	A	13/04/1999
				DE	69709075	D, T	22/08/2002
				DK	890013	T	08/04/2002
				EP	0890013	A, B	13/01/1999
				SE	0890013	T3	
				GB	9700862	D	00/00/0000
				NO	984548	A	27/11/1998
				US	6170578	B	09/01/2001

---